

ACCESS

Ďakujeme vám za vašu účasť na štúdiu **Access**.

Na nasledujúcich stranách uveďte údaje o pacientkách pozbierané v perioperačnom období cisárskeho rezu.

Budeme potrebovať, aby ste pre každú pacientku vyplnili celý dotazník. Všetky polia sú povinné.

Začnime!

Úvod

1 V ktorej krajine vykonávate svoju prax?

2 Aká je medzinárodná predvoľba vašej krajiny?

3 Číslo vášho zdravotníckeho zariadenia v rámci štúdie

4 Počet zdokumentovaných prípadov vo vašom centre

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12
- 13
- 14
- 15
- 16
- 17
- 18
- 19
- 20
- 21
- 22
- 23
- 24
- 25
- 26
- 27
- 28
- 29
- 30
- 31
- 32
- 33
- 34
- 35
- 36
- 37
- 38
- 39
- 40
- 41
- 42
- 43
- 44
- 45
- 46
- 47
- 48
- 49
- 50
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60
- 61
- 62
- 63
- 64
- 65
- 66
- 67
- 68
- 69

- 70
- 71
- 72
- 73
- 74
- 75
- 76
- 77
- 78
- 79
- 80

Podrobnosti o pacientovi

- 5 Veková skupina pacientky podstupujúcej Cisársky rez (SC) - (roky)?
- 15-20
 - 21-25
 - 26-30
 - 31-34
 - 35-39
 - 40-44
 - >44
-
- 6 Aká je výška (cm) pacientky podstupujúcej cisársky rez?
- _____
-
- 7 Aká je hmotnosť (kg) pacientky v čase cisárskeho rezu?
- _____
-
- 8 Aký je zdravotný stav pacientky (zaradte pacientku do ASA)?
- 1-2 Zdravá tehotná (obvykle klasifikovaná ASA 2) alebo Tehotná s nezávažným ochorením
 - 3 Tehotná so stredne závažným ochorením
 - 4 Tehotná so závažným ochorením
 - 5 Tehotná v život ohrozujúcej situácii
 - Iné: _____
-
- 9 Aká je najvyššia úroveň vzdelania/skúsenosti najskúsenejšieho anesteziológa prítomného v operačnej sále, ktorý vykonáva/dohliada na anestéziu tejto pacientky? (Supervízor nemusí byť prítomný v operačnej sále, ale je plne k dispozícii pomôcť.) Označte prosím najpresnejšiu možnosť.
- Anaesthesiologist trainee < 2 years
 - Trainee ≥ 2-5 years experience
 - Anaesthesiologist (trainee or attending) 6-10 years experience
 - Anaesthesiologist (trainee or attending) ≥10 years experience
 - Obstetric Anaesthesia specialist
 - Other: _____
-
- 10 Aká je indikácia k SC?
- Malpresentation
 - Multiple Gestation
 - Repeat Cesarean
 - Maternal Request
 - Placenta previa, Placenta accreta
 - Maternal medical condition
 - Intrapartum Cesarean emergency (eg failed trial of delivery/prolonged/ labour/ fair operative delivery)
 - Fetal compromise
 - Bleeding
 - Other: _____

- 11 Aká bola urgentnosť SC? Vyberte najviac vyhovujúcu možnosť.
- Emergentný SC/ červený kód: vybavenie plodu nutné do 15 minút
- Urgentný SC/oranžový kód: vybavenie plodu nutné do 30-60 minút
- Urýchlený SC/zelený kód: vybavenie plodu ím skôr od 60 minút do 24 hodín
- Plánovaný SC
-
- 12 Uveďte čas, kedy v priebehu dňa začal SC.
- 00:01 - 04:00
- 04:01 - 08:00
- 08:01 - 12:00
- 12:01 - 16:00
- 16:01 - 20:00
- 20:01 - 00:00
-
- 13 In case of unplanned Caesarean section, how long did it take from the call for Caesarean to fetal umbilical cord clamping (decision-to-delivery interval time,DDI) ? (minutes).
Please be as accurate as possible.
- Not Applicable
- Other: _____
-
- 14 What is the initial anaesthesia plan of the first anaesthesiologist to care for the patient (even if you then used another secondary plan)
- Spinal anaesthesia
- Epidural anaesthesia
- Combined spinal epidural
- Labor epidural top up
- Continuous spinal anaesthesia
- General anaesthesia
- other _____
-
- 15 Intrapartum Caesarean section: is there a labor epidural catheter in place for labor analgesia?
- Yes
- No
- Not Applicable
-
- 16 Intrapartum Cesarean section: In the case the patient has a labor epidural catheter in place, did the anaesthesiologist who made the initial anaesthesia mode plan decide/think that it was providing adequate labor analgesia?
- Not Applicable
- Yes
- No
-
- 17 Intrapartum Cesarean section: if a labour epidural was extended to provide anaesthesia, where was the top-up started?
- Not applicable
- In the labor room
- On the way to the operating room
- In the operating room
- other: _____

If neuraxial anaesthesia was used (Epidural top up, new epidural, new spinal anaesthesia or Combined Spinal Epidural Anaesthesia), please answer the questions below about local anaesthetic and adjuvants administered, with doses:

- 18 Spinal Local Anaesthetic Agent.
You may add drug dose/volume details to "Other".
- No spinal dose / not applicable
- Hyperbaric Bupivacaine 0.5%
- Isobaric Bupivacaine 0.5%
- Levobupivacaine 0.5%
- Chlorprocaine 3%
- Prilocaine 2%
- other: _____

-
- 19 Spinal lipophilic opioid. You may add drug dose/volume details to "Other".
- No spinal lipophilic opioid / not applicable
 Fentanyl
 Sufentanil
 other: _____
-
- 20 Spinal Long-Duration opioid. You may add drug dose/volume details to "Other".
- No spinal long-duration opioid / Not applicable
 Morphine
 Diamorphine
 other: _____
-
- 21 Spinal non opioid Adjuvants?
(Various options are possible)
- No spinal non-opioid adjuvant / Not applicable
 Clonidine
 Dexmedetomidine
 Epinephrine
 other: _____
-
- 22 Epidural Top-Up: Local Anaesthetic
You may add drug dose/volume details to "Other".
Please note that adjuvants are asked in a later question.
- No epidural top up / not applicable
 Ropivacaine 0.75%
 Ropivacaine 0.5%
 Levobupivacaine 0.5%
 Levobupivacaine 0.375%
 Bupivacaine 0.5%
 Bupivacaine 0.375%
 Lidocaine 2%
 Chloroprocaine 3%
 Other: _____
-
- 23 Epidural Top Up: Epidural lipophilic opioid. You may add drug dose/volume details to "Other".
- No epidural lipophilic opioid / not applicable
 Fentanyl
 Sufentanil
 Other: _____
-
- 24 Epidural Top Up: Epidural Long-duration Opioid.
You may add drug dose/volume details to "Other".
- No long-acting epidural opioid / not applicable
 Diamorphine
 Morphine
 Other: _____
-
- 25 Epidural Top Up: Non Opioid adjuvants (Various options are possible).
- No epidural non-Opioid adjuvant / not applicable
 Epinephrine
 Bicarbonate
 other: _____
-
- 26 Spinal / Epidural top up: Non opioid Adjuvants,
Comments: _____
-
- 27 If neuraxial anaesthesia was used (spinal/ topup of labor epidural analgesia/combined spinal-epidural), how was the block tested prior to incision?
Mark all applicable.
- Not tested
 Light touch test by the anaesthesiologist
 Pin-Prick test by the anaesthesiologist
 Cold test by the anaesthesiologist
 Motor Block test by the anaesthesiologist
 Skin test by surgeon
 other: _____

-
- 28 If the block was tested, what was the highest tested dermatome reached? (any testing modality)
- Not Tested
 T1 or above
 T2
 T3
 T4
 T5
 T6
 T7
 T8
 T9
 T10 or below
 Not applicable
-
- 29 If neuraxial anaesthesia was used (spinal/ topup of labor epidural analgesia/combined spinal-epidural), was there a failure of the block (requirement for repeat neuraxial procedure / conversion to general anesthesia) for Caesarean section?
- No
 Yes
 Not applicable
-
- 30 If yes, please indicate the type of neuraxial anesthesia failure (Various types can be selected).
- No Neuraxial Anaesthesia Failure / not applicable
 Complete failure of the block
 Low Level block
 Lateralized block
 Correct level but insufficient intensity of the block
 Other: _____
-
- 31 If there was a failure of neuraxial anaesthesia (spinal/ topup of labor epidural analgesia/ combined spinal-epidural) which of the following "next plan" strategies was used as a first option?
- No Neuraxial Anaesthesia Failure / not applicable
 General Anaesthesia
 Spinal Anaesthesia with usual dose of local anaesthetic
 Spinal Anaesthesia with reduced dose of local anaesthetic compared to routine practice
 Repeat epidural top up
 Combined Spinal-Epidural Anaesthesia
 Intravenous / Inhalational supplementation (clarifying - NOT general anaesthesia)
 Other: _____
-
- 32 If there was a failure of neuraxial anaesthesia (spinal/ topup of labor epidural analgesia/ combined spinal-epidural) which of the following "next plan" strategies was used as a second option?
- No second plan used/ not applicable
 General Anaesthesia
 Spinal Anaesthesia with usual dose of local anaesthetic
 Spinal Anaesthesia with reduced dose of local anaesthetic compared to routine practice
 Repeat epidural top up
 Combined Spinal-Epidural Anaesthesia
 Intravenous / Inhalational supplementation
 Other: _____
-
- 33 If there was a failure of neuraxial anaesthesia (spinal/ topup of labor epidural analgesia/ combined spinal-epidural) which of the following "next plan" strategies was used as a third option?
- No third plan used / not applicable
 General Anaesthesia
 Spinal Anaesthesia
 Repeat epidural top up
 Combined Spinal-Epidural Anaesthesia
 Intravenous / Inhalational supplementation
 Other: _____
-
- 34 Strategy for failed neuraxial block, comments (Optional):
- _____

-
- 35 If there was failure of neuraxial anaesthesia, when was the next plan (general anaesthesia/repeat block/intravenous/inhalational supplementation) initiated?
- No Neuraxial anaesthesia failure / not applicable
 - Before incision
 - After incision before delivery
 - After incision after delivery
-
- 36 If general anaesthesia (including invasive airway support) was performed during Caesarean section at any time, what was the reason?
- Maternal contraindication for neuraxial anaesthesia
 - Maternal request
 - Totally failed neuraxial anaesthesia
 - Partially failed neuraxial anaesthesia
 - Lack of time with insufficient level in a well-functioning block
 - Well-functioning block but patient preference to convert to GA
 - To support clinical management of haemodynamic instability
 - Not applicable
 - Other: _____
-
- 37 If intravenous / inhalational medications were administered as a supplement to neuraxial anaesthesia, please mark all used.
- No intravenous / inhalational medications
 - Propofol
 - Ketamine
 - Midazolam
 - Etomidate
 - Fentanyl / Alfentanil / Sufentanil / Remifentanil
 - Morphine
 - Nitrous Oxide
 - Sevoflurane
 - Other: _____